

Nutrazorb 45 is designed to be used as a foliar nutrient or soil applied as a starter with seeds or transplants.

APPLICATION:

May be applied by ground or air.

ALMONDS APRICOTS, CHERRIES, PLUMS PEACHES, NECTARINES: Apply at the rate of 1 gallon acre in sufficient water to obtain thorough coverage after bud swell to popcorn stage, prior to bloom. Repeat the application within two weeks after petal fall. Older trees may require another application of 3-4 quarts/acre 4-6 weeks later.

APPLES. PEARS: 2 to 4 quarts/acre in sufficient water prior to bloom. Repeat at petal fall. Another application applied 4-6 weeks water at the rate of 3-4 quarts/acre.

CAUTION: DO NOT APPLY TO MELROSE APPLES.

AVOCADOS: 3-4 quarts/acre in sufficient water prior to bloom. Two similar applications should be applied at 4-week intervals.

ALFALFA (Hay): 3-4 quarts/acre one week after each cutting.

ALFALFA (Seed): 3-4 quarts/acre at bud stage prior to bloom. 2-3 additional applications at the rate of 2-3 quarts/acre should be applied at 2-3 week intervals post bloom.

ARTICHOKES: 2-3 quarts/acre at bud stage. 2 quarts acre at 2 week intervals.

BARLEY, WHEAT, OTHER SMALL GRAINS: 2-3 quarts/acre prior to tillering. Another application of 2-3 quarts prior to heading out to help fill out kernels.

BEANS. PEAS: 2-3 quarts/acre at bud prior to bloom. Two additional applications at 2 week intervals, for set and size.

BUSH BERRIES, BOYSENBERRIES. BLACK-BERRIES, RASPBERRIES: 2-3 quarts/acre prior to bloom. 2-3 quarts/acre every 2 weeks thereafter.

CABBAGE, CAULIFLOWER, BROCCOLI, ALL COLE CROPS (Direct Seed): 2 quarts/acre at 4-5 true leaf stage. 2-3 quarts/acre at bud stage. 2-3 quarts/acre at two to three week intervals

CABBAGE, CAULIFLOWER, BROCCOLI, ALL COLE CROPS (Transplant): 1-2 quarts acre 1-2 weeks after transplanting. 2-3 quarts/acre at bud stage 2-3 quarts/acre at two to three week intervals.

CORN (Field & Sweet): 2-3 quarts/acre when corn is 2-4 feet tall. Repeat prior to tasseling.

SPINACH, CELERY: 2.3 quarts/acre after thinning lettuce and spinach. 1 week after transplanting celery. Repeat at 2-week intervals.

OTHER VEGETABLES & ROOT CROPS: 2-3 quarts/acre at 2-week intervals after thinning or when plants are well established.

CITRUS: 4.6 quarts acre in sufficient water for thorough coverage at bud stage prior to bloom. Repeat applications at 4-6 quarts/acre as new growth appears. Do not apply during bloom or just before harvest do not apply in combination with oils.

COTTON: 1-2 quarts/acre after true leaves appear. 2-3 quarts/acre at first squares. 2 additional applications at two week intervals.

GRAPES: 1-2 quarts/acre when canes reach 8-12 inches. *2-3 quarts/acre 1-2 weeks prior to bloom helps fruit set. 3 quarts acre between shatter and buckshot stage for sizing. 3 quarts/acre 4-6 weeks prior to harvest aids sugar content and color. *Omit second application for those varieties that set excessively or have tight clusters.

GROW MORE®



NUTRAZORB 45

8-32-5

Foliar or Soil

Total Nitrogen (N)	8%
8.0 Ammoniacal Nitrogen	
Available Phosphate (P2O5)	32%
Soluble Potash (K2O)	5%
Iron (Fe)	0.10%
0.10% Chelated Iron	
Zinc (Zn).....	0.05%
0.05% Chelated Zinc	

Plant nutrients derived from Ammonium , Tri Potassium Phosphate, Phosphoric Acid, Iron EDTA, Zinc EDTA and Humic Acid from Leonardite

MELONS, CUCUMBERS, SQUASH, CANTALOUPE: 2-3 quarts/acre at bud stage prior to bloom 2-3 additional applications at 2-week intervals.

PISTACHIO, PECANS, WALNUTS: 3-4 quarts/acre in sufficient water for thorough coverage before bloom. Repeat application at 3-4 quarts/acre every 6 weeks. The last application being applied 30 days before harvest.

IRISH POTATOES, SWEET POTATOES, YAMS: 3 quarts/acre when plants are 4-6 inches high. 2 additional applications at 3-4 week intervals.

RICE: 2-3 quarts/acre prior to tillering. Repeat 1-2 weeks prior to bloom. Repeat 2 weeks alter bloom.

CAUTION: DO NOT APPLY AT TIME OF BLOOM.

SAFFLOWER: 2-3 quarts/acre at bud stage prior to bloom. Repeat 2-3 weeks later.

STRAWBERRIES: 2-3 quarts/acre prior to bloom. Repeat at 2-week intervals throughout season.

SORGHUM: 2-3 quarts/acre prior to heading out. Repeat in 2-3 weeks.

SOYBEANS: 2-3 quarts/acre at bud prior to bloom. Repeat in 2 weeks. Apply an additional application of 2-3 quarts/acre as pods begin to fill.

SUGAR BEETS: 2-3 quarts/acre at thinning repeat 4 weeks later. Application of 2 to 3 quarts acre 6 weeks before harvest aids in increasing sugar content.

TOMATOES & PEPPERS (Processed): 1-2 quarts/acre after stand is established. 2-3 quarts/acre at bud stage prior to bloom. 2

additional applications at 2-week intervals.
TOMATOES & PEPPERS (Fresh): 2-3 quarts acre at bud stage prior to bloom. 2-3 quarts/acre 10-14 days throughout growing season. May be used in transplant water at the rate of 3-5 gallons/acre.

SHAKE OR STIR BEFORE USING.

NOTICE:

A wetting agent or spreader should be used when applying Nutrazorb 45 to prevent beading of droplets.

Buyer assumes all responsibility for safety and use not in accordance with the directions.

CAUTION:

KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED.

Avoid contact with skin, eyes or clothing.

In case of contact, immediately flush skin or eyes with plenty of water for at least 15 minutes.

WARNING: This product contains chemicals known to the State of California to cause cancer and birth defects and other reproductive harm.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

Manufactured By

GROW MORE®

15600 New Century Drive,
Gardena, CA 90248 U.S.A.

www.growmore.com

DENSITY: 11.7 LBS./GALLON

5 GALLONS (18.9 LITERS)