

OPTIMUM APPLICATION TIME: For best results, it is recommended that **Nutra-Zorb** be applied during the cool of the day, either in the early morning or late in the evening. Be sure equipment is clean and free of contamination.

NOTICE TO BUYER: Avoid prolonged exposure to dermis. Do not apply when weather conditions favor drip from the areas treated. Do not apply when runoff is likely to occur. Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur. Buyer assumes all risk of use and/or handling of this material, when such use and/or handling is contrary to label instructions.

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH CAUTION AND INSTRUCTIONS.

NOTICE: Seller's guaranty shall be limited to the terms of the label. The buyer assumes any risk to persons or property arising from use or handling and accepts the product on these conditions.

CAUTION: Causes eye irritation in case of contact with eyes, immediately flush with water for at least 15 minutes. If irritation persists, get medical attention. Do not swallow. Do not repack into unlabeled containers or those normally used for food or drink.

DENSITY: 10.4 Lbs./Gallon

**KEEP OUT OF REACH OF CHILDREN
CAUTION**



GROW MORE®

Research & Manufacturing Since 1918

15600 New Century Drive, Gardena, CA 90248

www.growmore.com

NUTRA-ZORB

8-8-8



GUARANTEED ANALYSIS:

Total Nitrogen (N).....	8.00%
2.5% Ammoniacal Nitrogen	
2.7% Nitrate Nitrogen	
2.8% Urea Nitrogen	
Available Phosphate (P ₂ O ₅).....	8.00%
Soluble Potash (K ₂ O).....	8.00%
Iron (Fe).....	0.10%
0.10% Chelated Iron	
Zinc (Zn).....	0.10%
0.10% Chelated Zinc	

ALSO CONTAINS NON-PLANT FOOD INGREDIENTS:

2% Organic acids derived from Leonardite

Derived from: Lo-biuret Urea, Ammonium Polyphosphate, Aqueous Ammonia, Potassium Hydroxide, Potassium Nitrate, Potassium Phosphate, Iron EDTA, Manganese Sulfate, and Zinc Citrate, Chelated with Citric Acid and EDTA (Ethylenediaminetetraacetic acid)

Grow More 8-8-8 is designed for application to agricultural crops including field and vegetable crops, vines, fruit trees and nut trees.

Application rates will vary according to the crop and cultivation practices.

Before using previously untried mixtures a jar test for compatibility is recommended. **8-8-8** may be foliar or soil applied by fertigation system or side dressed.

8-8-8 is tank mix compatible with most fungicides, herbicides and insecticides.

GENERAL USE:

FOLIAR: Use 1 to 12 quarts / acre depending on crop. Do not exceed concentration of 2 quarts in 10 gallons of water.

SOIL: Use 1 to 10 gallons / acre in fertigation system

GENERAL APPLICATION RATES AND TIMINGS:

Alfalfa (Foliar): 1 to 8 qt./acre. Apply after each hay cutting. On seed applications 4-6 times prior to harvest.

Artichokes (Foliar): 1 to 2 qt./acre. Apply at bud stage repeat beyond bud stage as needed.

Blackberries, Boysenberries, Raspberries, Strawberries (Foliar): 1 to 2 qt /acre. Apply at pre-bloom, repeat as needed until berries color or soften as varieties dictate

Broccoli, Brussel Sprouts, Cabbage, Cauliflower (Foliar): 1 to 2 qt./acre Apply at 2-leaf stage, repeat as needed.

Carrots (Foliar): 1 to 3 qt./acre. Apply when carrots are pencil-size, repeat as needed.

Cotton (Foliar): 1 to 8 qt./acre Apply at pre-bloom, repeat as needed.

Deciduous fruit and nut trees (Foliar): 1 to 12 qt./acre. Application schedule will be dependent on crop, variety and cultural practices of the region.

Garlic, Onions (Foliar): 1 to 2 qt./acre. Apply as needed.

Garlic, Onions (Fertigation/ Soil application): 1 to 10 gal./acre. Drench apply with irrigation water as needed.

Grapes (Foliar): 1 to 8 qt /acre. Apply after leaves emerge, use lower rates early. Increase rates as foliage increases.

Lettuce, Celery (Fertigation, Soil application): 1 to 10 gal./acre. Drench apply with irrigation water as needed.

Citrus, Avocados (Foliar): 1 to 12 qt./acre Application schedule will be dependent on crop, variety and cultural practices of the region.

Melons, Cucumbers, Squash (Foliar): 1 to 2 qt./acre. Apply when plant is 6-8" height, repeat as needed

Peas & Beans (Foliar): 1 to 2 qt /acre. Apply when plant is 6-8" height, repeat as needed.

Potatoes (Foliar): 1 to 8 qt. acre. Apply at bud formation; repeat as needed.

Soybeans (Foliar): 2 to 3 qt./acre. Apply at pre-bloom and after pod formation.

Corn (Foliar): 1 to 8 qt /acre. Apply at 6" height and repeat as needed.

Tomatoes, Eggplant, Peppers (Foliar): 1 to 2 qt./acre. Apply at first bloom, repeat after each fruit set.

Nursery, Greenhouse, Ornamentals (Drench): 12-36 oz./100 gal. Drench to soak planting medium.

Transplanting solutions (Drip or Drench): 12-36 oz./100 gal. Dip roots or drench planting medium.

**NET CONTENT: 2.5 GALLONS (9.46 LITERS)
NET WEIGHT: 26 LBS. (11.8 KG)**